Journal of Surgical Oncology

Table of Contents

Volume 90, Number 1, April 1, 2005

Review Article: Cryoablation and Cryolocalization in the Management of Breast Disease
Pat W. Whitworth and John C. Rewcastle 1
Cryotechnology has multiple applications in breast disease. Applications of cryoablation and cryolocalization will be discussed for use in diagnosing and treating benign breast conditions and breast cancer.

Review Article: Extremity Soft Tissue Sarcoma: Evidence-Based Multidisciplinary Management
Charles R. Scoggins and Raphael E. Pollock 10
Current management strategies for the treatment of primary extremity STS are outlined, with an emphasis on multidisciplinary care and protocol-based research.

Site-Select Procedure for Non-Palpable Breast Lesions: Feasibility Study With a 15-mm Cannula
Emmanuel Barranger, Olivier Marpeau, Jocelyne Chopier, Martine Antoine, and Serge Uzan 14
This study demonstrates that the Site-Select procedure is an effective diagnostic method in selected cases with low complication rate, high patient satisfaction, and excellent esthetic results.

Prognostic Significance of Circumferential Margin Involvement in Rectal Adenocarcinoma Treated With Preoperative Chemoradiotherapy and Low Anterior Resection
Pedro Luna-Pérez, Efraín Bustos-Cholico, Isabel Alvarado, Antonio Mafíuz, Saúl Rodríguez-Ramírez, Marcos Gutiérrez de la Barrera, and Sonia Labastida 20
Histologic examination of circumferential margins is an important predictor of local and distant relapse in non-radiated rectal cancer. However, for patients who received preoperative chemoradiotherapy this role has not yet been addressed. In patients with rectal cancer treated by preoperative chemoradiation plus total mesorectal excision and sphincter saving surgery, circumferential margin involvement is associated with high incidence of distant recurrence and cancer-related death.

Successful Use of Letrozole in Male Breast Cancer: A Case Report and Review of Hormonal Therapy for Male Breast Cancer
Brent P. Zabolotny, Christian V. Zalai, and Sarkis H. Meterissian 26

Spontaneous Rupture of Adrenal Pheochromocytoma: Review and Analysis of Prognostic Factors
Takashi Kobayashi, Akira Iwai, Ryo Takahashi, Yoshihiro Ide, Koji Nishizawa, and Kenji Mitsumori 31

Hepatectomy Plus Intraoperative Radiofrequency Ablation and Chemotherapy to Treat Technically Unresectable Multiple Colorectal Liver Metastases
Dominique Elias, Olivier Baton, Lucas Sideris, Valérie Boige, David Malka, Gabriel Liberale, Marc Pocard, and Philippe Lasser 36
The aim of this study was to assess the survival of patients with technically (and not oncologically) unresectable liver metastases when undergoing a combination of liver resection and radiofrequency with chemotherapy. This treatment lead to a median survival of 36 months and to a 2-year overall survival rate of 67%.

How I Do It: Bi-segmentectomy V-VIII as Alternative to Right Hepatectomy: An Intrahepatic Approach
Marcel Autran C. Machado, Paulo Herman, Roberto F. Meirelles Jr, Telesforo Bacchella, and Marcel C.C. Machado 43

Volume 90, Number 2, May 1, 2005

Guest Editorial: Distinctive Biology of Pleomorphic Lobular Carcinoma of the Breast
Roger E. Moe and Benjamin O. Anderson 47

Cytoreductive Surgery and Intraoperative Hyperthermic Chemoperfusion for Advanced Ovarian Carcinoma
Trevor W. Reichman, Bernadette Cracchiolo, Jahir Sama, Margarette Bryan, Jonathan Harrison, Lillian Pliner, and Lawrence E. Harrison 51
The use of intraoperative hyperthermic chemoperfusion (IHCP) coupled with optimal cytoreduction is a safe and effective treatment for advanced ovarian carcinoma. However, the proper selection of patients who will benefit most from the therapy is essential for the success of the treatment.

Commentary
Paul H. Sugarbaker 56

Detection of Peritoneal Micrometastasis by Reverse Transcriptase-Polymerase Chain Reaction for Heparanase mRNA and Cytology in Peritoneal Wash Samples
Zhenning Wang, Xue Zhang, Huimian Xu, Xin Zhou, Li Jiang, and Chong Lu 59
In this study, we have reported the clinical relevance of a new diagnostic method with RT-PCR assay using heparanase as a target gene for detection of free cancer cells in peritoneal washes. It is suggested that heparanase could be ubiquitously detected in peritoneal free cancer cells and metastatic nodules, and the high sensitivity of RT-PCR based heparanase assay made it a candidate molecular marker for detecting peritoneal micrometastasis.

Significance of Serum S-100B in Melanoma Patients Before and After Sentinel Node Biopsy
Léonie H.M. Smit, Omgo E. Nieweg, Catharina M. Korse, Johannes M.G. Bonfrer, and Bin B.R. Kroon 66
S-100B is not useful in predicting the presence of metastatic disease in the sentinel node. The biomarker has limited sensitivity but is highly specific in pointing out previously unknown metastases during follow-up.

Commentary
Mark B. Faries, Rishab K. Gupta, and Donald L. Morton 69

Ductal Carcinoma In Situ in Core Biopsies Containing Invasive Breast Cancer: Correlation With Extensive Intraductal Component And Lumpectomy Margins
Martin Dzierzanowski, Karen A. Melville, Penny J. Barnes, Rebecca F. MacIntosh, Judy S. Caines, and Geoffrey A. Porter 71
Until now, the significance of ductal carcinoma in situ (DCIS) found in core biopsies (CB) containing invasive breast cancer was unclear. The objective of this study was to examine the implications of DCIS found in the original CB, specifically related to the risk of extensive intraductal component and/or positive lumpectomy margins.

Quantification of Angiogenesis Induced in Rabbit Cornea by Breast Carcinoma of Women Treated With Tamoxifen
Benedito Borges da Silva, Raimundo Gerônimo da Silva Júnior, Umbelina Soares Borges, Marcos Antônio Guedes da Silveira Filho, Igor Clausius Carvalho Pimentel, Luiz Henrique Gebrim, Manuel de Jesus Simões, and Edmund Chada Baracat 77
Tamoxifen inhibited angiogenesis induced by breast carcinoma in rabbit cornea.

Activation of MMP-2 and MMP-9 in Patients With Oral Squamous Cell Carcinoma
Beena P. Patel, Pankaj M. Shah, Upendra M. Rawal, Amisha A. Desai, Shakuntala V. Shah, Rakesh M. Rawal, and Prabhudas S. Patel 81

Expression of RCAS1 in Esophageal Squamous Cell Carcinoma Is Associated With a Poor Prognosis
Hiroyuki Kato, Masanobu Nakajima, Norihiro Masuda, Ahmad Faried, Makoto Sohda, Yasuyuki Fukai, Tatsuya Miyazaki, Minoru Fukuchi, Katsuhiko Tsukada, and Hiroyuki Kuwano 89

Extended Multiorgan Resection for T4 Gastric Carcinoma: 25-Year Experience
Fabio Carboni, Pasquale Lepiane, Roberto Santoro, Riccardo Lorusso, Pietro Mancini, Isabella Sperduti, Massimo Carlini, and Eugenio Santoro 95
Surgical management of locally advanced gastric carcinoma invading adjacent organs remains controversial. However, the survival of patients can be improved with extended multiorgan resection, with acceptable postoperative morbidity and mortality rates, provided that a curative resection is performed.

Case Report: Extended Survival in a Patient With Recurrent and Metastatic Adrenal Cortical Carcinoma by Aggressive Transarterial Embolization—a Case Report
Shau-Hsuan Li, Cheng-Hua Huang, Sheung-Fat Ko, Fong-Fu Chou, and Shun-Chen Huang 101

Erratum 106

Volume 90, Number 3, June 1, 2005

Seminars in Surgical Oncology: Gastric Cancer
Walter Lawrence, Jr., MD, Guest Editor

Intro: Walter Lawrence, Jr.: A Tribute to a Surgical Oncologist. "Been There, Done That"
Harry D. Bear 109

Guest Editor Comment: Gastric Cancer: An Update
Walter Lawrence, Jr. 113

Gastric Cancer: New Genetic Developments
Henry T. Lynch, William Grady, Gianpaolo Suriano, and David Huntsman 114

COMMENT
Walter Lawrence, Jr. 133

New Strategies for the Prevention of Gastric Cancer: *Helicobacter pylori* and Genetic Susceptibility
Pelayo Correa 134

COMMENT
Walter Lawrence, Jr. 138

Biologic and Clinical Variations of Adenocarcinoma at the Esophago–Gastric Junction: Relevance of a Topographic-Anatomic Subclassification
J. Rüdiger Siewert, Marcus Feith, and Hubert J. Stein 139

COMMENT
Walter Lawrence, Jr. 146

Recent Advances in Sentinel Node Navigation for Gastric Cancer: A Paradigm Shift of Surgical Management
Yuko Kitagawa, Hirofumi Fujii, Koichiro Kumai, Tetsuro Kubota, Yoshihide Otani, Yoshiro Saikawa, Masashi Yoshida, Atsushi Kubo, and Masaki Kitajima 147

COMMENT
Walter Lawrence, Jr. 151

Status of Extended Lymph Node Dissection: Locoregional Control Is the Only Way to Survive Gastric Cancer
Henk H. Hartgrink and Cornelis J.H. van de Velde 153

Role of Post-Operative Chemoradiation in Resected Gastric Cancer
John S. Macdonald 166

Invited Commentary: Eliminating and Suppressing Local-Regional Disease in Gastric Cancer
Scott A. Hundahl 171

COMMENT
Walter Lawrence, Jr. 173

Perioperative Adjunctive Treatment in the Management of Operable Gastric Cancer
Howard Silberman 174

COMMENT
Walter Lawrence, Jr. 186

Minimally Invasive Treatment for Gastric Cancer: Approaches and Selection Process
Sung Hoon Noh, Woo Jin Hyung, and Jae-Ho Cheong 188

COMMENT
Walter Lawrence, Jr. 193

Gastric GI Stromal Tumors (GISTs): The Role of Surgery in The Era of Targeted Therapy
Michael C. Heinrich and Christopher L. Corless 195

COMMENT
Walter Lawrence, Jr. 207

Volume 90, Number 4, June 15, 2005

Prospective Non-Randomized Trial Comparing Esophagectomy-Followed-By-Chemoradiotherapy Versus Chemoradiotherapy-Followed-By-Esophagectomy for T4 Esophageal Cancers
Hirmasa Fujita, Susumu Sueyoshi, Toshiaki Tanaka, Yuichi Tanaka, Hiroko Sasahara, Kazuo Shirouzu, Gen Suzuki, Naofumi Hayabuchi, and Hiroki Inutsuka 209
Forty-three patients with a T4 esophageal cancer were enrolled in a prospective non-randomized study in which each patient decided on which treatment arm to receive: chemoradiotherapy-followed-by-esophagectomy or esophagectomy-followed-by-chemoradiotherapy. This trial concluded that the first choice of treatment for such patients was prior chemoradiotherapy rather than prior esophagectomy.

Impact of the Extent and Duration of Cytoreductive Surgery on Postoperative Hematological Toxicity After Intraperitoneal Chemohyperthermia for Peritoneal Carcinomatosis
Dominique Elias, Bruno Raynard, Valérie Boige, Agnes Laplanche, George Estphan, David Malka, and Marc Pocard 220

Surgical Results for Hepatocellular Carcinoma With Bile Duct Invasion: A Clinicopathologic Comparison Between Macroscopic and Microscopic Tumor Thrombus
Minoru Esaki, Kazuaki Shimada, Tsuyoshi Sano, Yoshihiro Sakamoto, Tomoo Kosuge, and Hidenori Ojima 226

Expression of L-Type Amino Acid Transporter 1 (LAT1) in Esophageal Carcinoma
Hideaki Kobayashi, Yukimoto Ishii, and Tadatoshi Takayama 233

Predictive Value of Lymph Node and Tumor Matrix Metalloproteinase Expression in the Analysis of Metastatic Periampullary Tumors
Gregory J. McKenna, Adam Meneghetti, Yong Liang Chen, Alice L.F. Mui, Christopher Ong, Charles H. Scudamore, W. Robert McMaster, David A. Owen, and Stephen W. Chung 239
16 patients with periampullary malignancies were followed prospectively to determine the whether there was a correlation between expression of matrix metalloproteinases and clinical outcomes.

How I Do It: A Modified Technique for Insertion of Intraperitoneal Port for Chemotherapy
Wen-Shiung Liou, Nelson N.H. Teng, and John K. Chan 247

Review Article: Apoptotic and Anti-Angiogenic Strategies in Liver and Gastrointestinal Malignancies
Jannis Kountouras, Christos Zavos, and Dimitrios Chatzopoulos 249
Inducing the apoptotic process in various ways and targeting fundamental play-markers of the angiogenic process can be applied as novel strategies to cancer management.

Author Index to Volume 90 260

Subject Index to Volume 90 262